Scientific Article



Title

Misophonia: diagnostic criteria for a new psychiatric disorder

Language: EN, Dutch

Authors

Arjan Schroeder (1), Nienke Vulink (1); Damiaan Denys (1-2)

- 1 Department of Psychiatry, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands
- 2. The Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Amsterdam, The Netherlands

Abstract

42 patients who reported misophonia were recruited by our hospital website. All patients were interviewed by an experienced psychiatrist and were screened with an adapted version of the Y-BOCS, HAM-D, HAM-A, SCL-90 and SCID II. The misophonia patients shared a similar pattern of symptoms in which an auditory or visual stimulus provoked an immediate aversive physical reaction with anger, disgust and impulsive aggression. The intensity of these emotions caused subsequent obsessions with the cue, avoidance and social dysfunctioning with intense suffering. The symptoms cannot be classified in the current nosological DSM-IV TR or ICD-10 systems.

The authors suggest that misophonia should be classified as a discrete psychiatric disorder. Diagnostic criteria could help to officially recognize the patients and the disorder, improve its identification by professional health carers, and encourage scientific research.

Source

PubMed, Journal List, PMID: 23372758. Published in PlosObe 2013; 8(1): e54706. online 2013 Jan 23.



Project number: 2021-1-BE02-KA220-VET-000034849

Scientific Article



Link

Retrieved from Plos One, Global Public Health, January, 2023: 10.1371/journal.pone.0054706 https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0054706

