

# Scientific Article



## Title

The Brain Basis for Misophonia

Language: EN

## Authors

Sukhbinder Kumar,<sup>1,2,7,\*</sup> Olana Tansley-Hancock<sup>1</sup>, William Sedley<sup>1</sup>, Joel S. Winston<sup>2,3</sup>, Martina F. Callaghan<sup>2</sup>, Micah Allen<sup>2</sup>, Thomas E. Cope<sup>1,4</sup>, Phillip E. Gander<sup>5</sup>, Doris-Eva Bamiou<sup>6</sup>, and Timothy D. Griffiths<sup>1,2</sup>

1 Institute of Neuroscience, Medical School, Newcastle University, Newcastle upon Tyne NE2 4HH, UK

2 Wellcome Trust Centre for Neuroimaging, 12 Queen Square, London WC1N 3BG, UK

3 Institute of Cognitive Neuroscience, 17 Queen Square, London WC1N 3AR, UK

4 Department of Clinical Neurosciences, University of Cambridge, Cambridge CB2 0SZ, UK

5 Human Brain Research Laboratory, Department of Neurosurgery, The University of Iowa, Iowa City, IA 52242, US

6 UCL Ear Institute, 332 Grays Inn Road, London WC1X 8EE, UK

7 Lead contact

\* Correspondence by S. Kumar

## Abstract

Using functional and structural MRI coupled with physiological measurements, we demonstrate that misophonic subjects show specific trigger-sound-related responses in brain and body. Specifically, fMRI showed that in misophonic subjects, trigger sounds elicit greatly exaggerated blood-oxygen-level-dependent (BOLD) responses in the anterior insular cortex (AIC), a core hub of the “salience network” that is critical for perception of interoceptive signals and emotion processing. Trigger sounds in misophonics were associated with abnormal functional connectivity between AIC and a network of regions responsible for the processing and regulation of emotions, including ventromedial prefrontal cortex (vmPFC), posteromedial cortex (PMC), hippocampus, and amygdala. Trigger sounds elicited heightened heart rate (HR) and galvanic skin response (GSR) in misophonic subjects, which were mediated by AIC activity. Questionnaire analysis showed that misophonic subjects perceived their bodies differently: they scored higher on interoceptive sensibility than controls, consistent with abnormal functioning of AIC. Finally, brain structural measurements implied greater myelination within vmPFC in misophonic individuals. Overall, our results show that misophonia is a



Co-funded by the  
Erasmus+ Programme  
of the European Union

Project number: 2021-1-BE02-KA220-VET-000034849

# Scientific Article



disorder in which abnormal salience is attributed to particular sounds based on the abnormal activation and functional connectivity of AIC.

## Source

ELSEVIER, Cell Press:

- Kumar et al., 2017, Current Biology 27, 527–533
- February 20, 2017 © 2016 The Author(s). Published by Elsevier Ltd.

## Link

Retrieved January 2023 from <http://dx.doi.org/10.1016/j.cub.2016.12.048>



Co-funded by the  
Erasmus+ Programme  
of the European Union

Project number: 2021-1-BE02-KA220-VET-000034849